Earth's Climate: Past, Present and Future

Thursday March 5th, 2015

Paul Belanger

Earth's past climate history

1. Earth's deep past before the Cambrian (600 MaBP): hot and cold
2. Earth's past: Cambrian onward: mostly hot-house Earth; 100s parts per million (ppm)
3. Climate trend in the Cenozoic – the last 65 million years; proxy data from 3600ppm to <200 ppm.
4. More recent past: 180-280 part per million; how do we know – empirical data. Preview of next week's field trip
5. Today: 400 ppm and growing 2-3ppm/year

climate system - the inter-relationship and feedback of:

- **Atmosphere**
- **Hydrosphere**
- **Biosphere**
- **Cryosphere**
- **Lithosphere (weathering reduces CO_2; volcanism increases it)**

INTRODUCTION: Definitions:

- **First order Forcings:** EXTERNAL Influences (3):

SOLAR input:



0.9% less
100 My ago



Atmospheric Opacity
(gases that absorb radiation in or out)

Albedo (30-85%)

- **Feedbacks:** INTERNAL dynamics and responses
  - e.g. higher water vapor in atm. due to heating of atm



Figure 2.11: | Global mean energy budget under present-day climate conditions. Numbers state magnitudes of the individual energy fluxes in W m^{-2}, adjusted within their uncertainty ranges to close the energy budgets. Numbers in parentheses attached to the energy fluxes cover the range of values in line with observational constraints. (Adapted from Wild et al., 2013.)

161+342 = 503 - 2 outside
vs. 84+20+398=502 - 3 inside
arrows

84 +20 +56 = 160
which =~ incoming 161 shortwave



10°C = (50°F)	20°C = (68°F)	30°C = (86°F)	40°C = (104°F)
7.8 cc	15 cc	27.7 cc	49.8 cc

The Earth and its atmosphere

Water vapor

CO_2

The most potent greenhouse gas is H_2O - vapor

The large H_2O greenhouse effect is controlled by temperature – H_2O saturation doubles with every 10°C Increase

As a result It is concentrated in the lower atmosphere of the tropics

The CO_2 greenhouse gas effect is concentrated in the polar regions ! ! !

Particularly in the Arctic !

CO_2 is evenly distributed throughout the atmosphere, there is little water at the poles because cold air holds less H_2O

Earth's past climate

1. Earth's deep past before the Cambrian (600 MaBP): hot and cold
2. Earth's past: Cambrian onward: mostly hot-house Earth; 100s parts per million (ppm)
3. Climate trend in the Cenozoic – the last 65 million years; proxy data from 3600ppm to <200 ppm.
4. More recent past: 180-280 part per million; how do we know – empirical data. Preview of next week's field trip
5. Today: 400 ppm and growing

Earth's past climate

Earth's deep past and early atmosphere before the Cambrian (600 MaBP): hot and cold

- Earth self regulates 2.1 -2.3 Tim Lenton video – 9 minute overview
- Article Link: BBC Nature http://www.bbc.co.uk/nature/ancient_earth/Snowball_Earth
- You Tube – leaving for you to watch on your own:
- https://www.youtube.com/watch?v=mX3pHD7NH58
- https://www.youtube.com/results?search_query=snow+ball+earth – various links

Earth's past climate

1. Earth's deep past before the Cambrian (600 MaBP): hot and cold
2. **Earth's past: Cambrian onward: mostly hot-house Earth; 100s parts per million (ppm)**
3. Climate trend in the Cenozoic – the last 65 million years; proxy data from 3600ppm to <200 ppm.
4. More recent past: 180-280 part per million; how do we know – empirical data. Preview of next week's field trip
5. Today: 400 ppm and growing


Phanerozoic Climate Change
Cm
O
S
D
C
P
Tr
J
K
Pg
N
Short-Term Average
Long-Term Average
HOT
COLD
Glacial Periods
-4
-3
-2
-1
0
1
2
3
δ18O (parts per thousand)
542
500
450
400
350
300
250
200
150
100
50
0
Millions of Years Ago

Climate Changes from Ocean Sediment Cores, since 5 Ma. Milankovitch Cycles



When CO_2 levels get below ~400-600 ppm Orbital parameters become more important than CO_2

Earth's past climate

1. Earth's deep past before the Cambrian (600 MaBP): hot and cold
2. Earth's past: Cambrian onward: mostly hot-house Earth; 100s parts per million (ppm)
3. **Climate trend in the Cenozoic – the last 65 million years; proxy data from 3600ppm to <200 ppm.**
4. More recent past: 180-280 part per million; how do we know – empirical data. Preview of next week's field trip
5. Today: 400 ppm and growing

Scientific History of Climate change – PROXY DATA







Alternating Greenhouse Earth / Ice-house Earth

Geologic cycles:
Climate through the Phanerozoic:
Carbon is the culprit



Royer et al., 2003

Cenozoic Deep Sea Climate Record



Zachos et al. 2008

Correlation of CO_2 and temperature over last 65 million years



Beerling and Royer, Nature 2011

Long-term Carbon Cycle: rocks

Two generalized reactions…

Photosynthesis/Respiration

$$CO_2 + H_20 \leftrightarrow CH_2O + O_2$$

Weathering/Precipitation

$$CO_2 + CaSiO_3 \leftrightarrow CaCO_3 + SiO_2$$

Silicate weathering

Regulates atmospheric CO_2 on geologic time scales.



J. Ormes, 2015

Long-term carbon cycle: *rocks*



Berner, 2001

^{18}O as a temperature proxy

Evaporation and condensation influence the ratio of $^{18}O/^{16}O$.
H_2O containing ^{16}O evaporates slightly more readily than H_2O containing ^{18}O.
H_2O containing ^{18}O condenses slightly more readily than H_2O containing ^{16}O.

Heavy hydrogen (deuterium) can be used in the same way.



J. Ormes, 2015



50 million years ago (50 MYA) Earth was ice-free.

Atmospheric CO_2 amount was of the order of 1000 ppm 50 MYA.

Atmospheric CO_2 imbalance due to plate tectonics ~ 10^{-4} ppm per year.


Global Deep Ocean Temperature
PETM
Antarctic Ice Sheet
N. Hemisphere Ice Sheets
India Collides
with Eurasia
Antarctic
Glaciation
Temperature (°C)
12
8
4
0
60
50
40
30
20
10
0
Time (Million Years Before Present)


Antarctic ice sheet
12
8
4
0
Pal.
Eocene
Olig.
Miocene
Plio.
2,000
1,500
1,000
500
0
Stomata
Phytoplankton
Palaeosols
Liverworts
Boron
B/Ca
Nahcolite–trona
60
50
40
30
20
10
0
Time (Myr ago)

Earth's past climate

1. Earth's deep past before the Cambrian (600 MaBP): hot and cold
2. Earth's past: Cambrian onward: mostly hot-house Earth; 100s parts per million (ppm)
3. Climate trend in the Cenozoic – the last 65 million years; proxy data from 3600ppm to <200 ppm.
4. **More recent past: 180-280 part per million; how do we know – empirical data. Preview of next week's field trip**
5. Today: 400 ppm and growing

Climate Changes from Ocean Sediment Cores, since 5 Ma. Milankovitch Cycles



When CO_2 levels get below ~400-600 ppm Orbital parameters become more important than CO_2

Earth's past climate

1. Earth's deep past before the Cambrian (600 MaBP): hot and cold
2. Earth's past: Cambrian onward: mostly hot-house Earth; 100s parts per million (ppm)
3. Climate trend in the Cenozoic – the last 65 million years; proxy data from 3600ppm to <200 ppm.
4. More recent past: 180-280 part per million; how do we know – empirical data. Preview of next week's field trip
5. **Today: 400 ppm and growing**

- SO –
WHAT CONTROLS CLIMATE



Gerhard et al., 2001

FEEDBACKS



Rohling, et al., (PALAESENS Project mbrs), 2012

Paleocene/Eocene Thermal Maximum
PETM


Global Deep Ocean Temperature
PETM
Antarctic Ice Sheet
N. Hemisphere Ice Sheets
India Collides
with Eurasia
Antarctic
Glaciation
Temperature (°C)
12
8
4
0
60
50
40
30
20
10
0
Time (Million Years Before Present)


Antarctic ice sheet
12
8
4
0
Pal.
Eocene
Olig.
Miocene
Plio.
2,000
1,500
1,000
500
0
Stomata
Phytoplankton
Palaeosols
Liverworts
Boron
B/Ca
Nahcolite–trona
60
50
40
30
20
10
0
Time (Myr ago)

Proxy data: stable isotopes



Wikipedia


4 45 46 47 48 49 50 51 52 4 55 56 5
PMAG


1263
2717 m water depth
1265
3060 m water depth
1266
3798 m water depth
1267
4355 m water depth
1262
4755 m water depth
Depth (mcd)
334
335
336
315
316
305
306
307
230
231
232
139
140
BF-4
BF-3
BF-2
BF-1
100 50 0
CaCO3 (wt%)

PETM - THE LAND RECORD

Bighorn Basin

- PETM interval in fluvial deposits with excellent alluvial paleosols - seen as color bands, which are soil horizons
- Found in Willwood Fm
- Reds, purples due to iron oxides in B horizons

Paleosol Density

Bighorn Basin Climate

- Plant fossils and isotopes show Mean Annual Temperature of 20° to 25° C or 68 to 77° F

- Similar to Gulf Coast region today



PROXY DATA-EXTRAS



FROM CSI TO GSI: GEOLOGICAL SAMPLE INVESTIGATION

LET THE EVIDENCE SPEAK FOR ITSELF



WE CALL THIS EVIDENCE

"PROXY" DATA

SOME OF THE EARLIEST PROXY DATA WAS FROM TERRESTRIAL DEPOSITS

- Strandlines/shorelines
- Moraines
- Till
- Kettle lakes, etc.

We may know what caused these today, but imagine back then?

IT'S THE INTERPRETATION THAT'S NOT ALWAYS CORRECT

Darwin observed ancient Alpine shorelines:
interpreted as ocean shoreline

Agassiz – later correctly interpreted as ice-dammed lake-shore strandlines/shoreline

Louis Agassiz

Louis Agassiz

Born	May 28, 1807 Haut-Vully, Switzerland
Died	December 14, 1873 (aged 66) Cambridge, Massachusetts
Fields	Paleontology, Glaciology, Geology, Natural History
Alma mater	University of Erlangen-Nuremberg

- **Jean Louis R. Agassiz**
- **"Father" of Glaciology**
- 1807-1873
- Paleontologist
- Glaciologist

Photographic proxy data/evidence



Ruddiman, 2008

EARLY PROXY DATA: TREE RINGS



Pollen & Lake core data



Ruddiman, 2008



PROXY DATA: POLLEN DATA

PROXY DATA: LEAVES

Tree rings, corals, ice cores



Ruddiman, 2008

PROXY DATA: ICE CORES



TERRESTRIAL DATA

North American:
- Wisconsin
- Illinoian
- Kansan
- Nebraskan

European:
- Wurm
- Riss
- Mindel
- Gunz

LATER EVIDENCE CAME FROM THE MARINE RECORD

NOT WITHOUT IT'S PROBLEMS, BUT MORE COMPLETE

Cesare Emiliani in the early 1950s when he was doing his pioneering research at the University of Chicago (Photo from the Archives of the Rosenstiel School of Marine and Atmospheric Science, University of Miami).

Cesare Emilani:

Paleontologist, Chemist

Father of Paleoceanography

Other Paleoceanographers

Wally Broecker

Thermal-haline

"conveyor" belt of circulation



Other Paleoceanographers

Bill Ruddiman





Nick Shackleton



Other Paleoceanographers

John Imbrie:
CLIMAP









PROXY DATA: CORE DATA

PROXY DATA: BENTHIC FORAMS









PROXY DATA: PLANKTONIC FORAMS

Deep Sea Coring



Ruddiman, 2008

A Climate knowledge quiz:

- http://www.csmonitor.com/Environment/2014/0827/Climate-change-Is-your-opinion-informed-by-science-Take-our-quiz/Gas

N.B. Wait until the end of the class to take this quiz. Jonathan F. Ormes

IS CLIMATE CHANGE ALL DOOM AND GLOOM?

- **Answer: NO!**
- **– but YES it's a challenge and as humans we have always been challenged: read the history books.**
- **That doesn't mean we stick our heads in the sand and ignore the challenges.**
- **We have human ingenuity and adaptability.**
- **Is ignoring and doing nothing an appropriate answer.**
- **Is resigning oneself to abdicating addressing the issue merely a way of dismissing and saying there's nothing we can do.**
- **There are a lot of smart people working to meet those challenges – let me cite 2 that might be "game changers".**
- More at: http://denverclimatestudygroup.com/